

(19) World Intellectual Property Organization  
International Bureau



(43) International publication date  
1 September 2005 (01.09.2005)

PCT

(10) International publication number  
WO 2005/080312 A1

(51) International patent classification<sup>7</sup>:  
C07C 69/716, D06M 13/188

(21) International application number: PCT/FR2005/000123

(22) International filing date: 20 January 2005 (20.01.2005)

(25) Language of filing: French

(26) Language of publication: French

(30) Data relating to the priority:  
04/00,666 23 January 2004 (23.01.2004) FR

(71) Applicant (for all designated States except US): ISOICHEM  
[FR/FR]; 9, rue Lavoisier, F-91170 Vert-le-Petit (FR).

(72) Inventors; and

(75) Inventors/Applicants (US only): FERRUCIO, Laurence  
[FR/FR]; 30, Domaine du Guichet, F-91810 Vert le Grand  
(FR). IERNO, Hélène [FR/FR]; 6, rue Maurice Boyau, F-  
91220 Bretigny sur Orge (FR).

(74) Representatives: MARTIN, Jean-Jacques, etc.; Cabinet  
Regimbeau, 20, rue de Chazelles, F-75847 Paris Cedex 17  
(FR).

(81) Designated states (unless otherwise indicated, for every kind  
of national protection available): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,

CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB,  
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,  
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ,  
OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,  
SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(81) Designated states (unless otherwise indicated, for  
every kind of regional protection available): ARIPO  
(BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ,  
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ,  
MD, RU, TJ, TM), European (AT, BE, BG, CH, CY,  
CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT,  
LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,  
MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

- Of inventorship (Rule 4.17(iv)) for the following  
designation US.

Published:


- With International Search Report.

For an explanation of the two-letter codes and the other  
abbreviations, reference is made to the explanations  
("Guidance Notes on Codes and Abbreviations") at the  
beginning of each regular edition of the PCT Gazette.

As printed

(54) Title: ACTIVE UNSATURATED ESTERS, METHOD FOR THE PRODUCTION THEREOF, AND USE OF THE SAME

(54) Titre : ESTERS INSATURES ACTIVES, PROCEDE DE FABRICATION ET UTILISATION



(I), according to which the compound of formula (II) is formed by the reaction between an alcohol and an acid anhydride, and by the activation of the carbonyl function. Furthermore, the invention relates to the use of the compound of formula (I) for increasing the hydrophobic character of polymers comprising amine and/or hydroxyl functions by means of grafting, and to a support based on natural textile fibres onto which the compound of formula (I) is grafted.

(57) Abrégé : La présente invention concerne un composé de formule (I) dans laquelle X représente un groupement activant, tel qu'un atome de chlore ou que l'imidazole, Y représente un radical aliphatique en C<sub>6</sub>-C<sub>20</sub> saturé et Z représente un radical aliphatique en C<sub>2</sub>-C<sub>10</sub>. La présente invention concerne également un procédé de synthèse du composé de formule (I) consistant en la formation du composé de formule (II) par réaction entre un alcool et un anhydride d'acide, puis activation de la fonction carbonyle. La présente invention concerne enfin l'utilisation du composé de formule (I) pour augmenter le caractère hydrophobe de polymères comportant des fonctions amines et/ou hydroxyles par greffage ainsi qu'un support à base de fibres textiles naturelles sur lequel est greffé le composé de formule (I).

(57) Abstract: The invention relates to a compound of formula (I) wherein X represents an activating radical such as a chlorine atom or imidazole, Y represents a saturated aliphatic C<sub>6</sub>-C<sub>20</sub> radical, and Z represents an aliphatic C<sub>2</sub>-C<sub>10</sub> radical. The invention also relates to a method for synthesising the compound of formula (I) consisting in the formation of the compound of formula (II) by reaction between an alcohol and an acid anhydride, and by activation of the carbonyl function. Furthermore, the invention relates to the use of the compound of formula (I) for increasing the hydrophobic character of polymers comprising amine and/or hydroxyl functions by means of grafting, and to a support based on natural textile fibres onto which the compound of formula (I) is grafted.